

Series representations of the remainders in the expansions for certain trigonometric functions and some related inequalities, II

Chao-Ping Chen
Richard B. Paris

This version of the article has been accepted for publication, after peer review and is subject to [Springer Nature's AM terms of use](#), but is not the Version of Record and does not reflect post-acceptance improvements, or any corrections. The Version of Record is available online at: <http://dx.doi.org/10.1007/s13398-022-01208-6>

Series representations of the remainders in the expansions for certain trigonometric functions and some related inequalities. II

C.-P. CHEN^a AND R. B. PARIS^b

^a *School of Mathematics and Informatics, Henan Polytechnic University,
Jiaozuo City 454000, Henan Province, China*

E-Mail: chenchaoping@sohu.com

^b *Division of Computing and Mathematics, University of Abertay Dundee,
Dundee DD1 1HG, UK*

E-Mail: r.paris@abertay.ac.uk

Abstract

We examine Wilker and Huygens-type inequalities involving trigonometric functions making use of results derived in Part I. The Papenfuss-Bach inequality representing upper and lower bounds for the function $x \sec^2 x - \tan x$ for $0 \leq x < \frac{1}{2}\pi$ is also investigated. An open problem posed by Sun and Zhu concerning this last inequality is established.

MSC: 11B68, 25D05

Keywords: Wilker inequality, Huygens inequality, Papenfuss-Bach inequality, trigonometric functions, inequalities

1. Introduction

In Part I [6] we derived series representations of the remainders in the expansions of certain trigonometric and hyperbolic functions and discussed some inequalities involving these functions. In Part II we apply some of the results obtained in [6] in the discussion of further trigonometric inequalities that are related to the Wilker, Huygens and Papenfuss-Bach inequalities.

In [24], Wilker proposed the following two open problems: Show that if $0 < x < \frac{1}{2}\pi$, then

$$\left(\frac{\sin x}{x}\right)^2 + \frac{\tan x}{x} > 2 \quad (1.1)$$

and also find the largest constant c when the right-hand side of (1.1) is replaced by $2 + cx^3 \tan x$. This was proved in [22] and further extended by Chen and Cheung in [5], where it was shown that for $0 < x < \frac{1}{2}\pi$

$$2 + \frac{8}{45}x^4 + \frac{16}{315}x^5 \tan x < \left(\frac{\sin x}{x}\right)^2 + \frac{\tan x}{x} < 2 + \frac{8}{45}x^4 + \left(\frac{2}{\pi}\right)^6 x^5 \tan x$$

with the constants $\frac{16}{315}$ and $(2/\pi)^6$ being best possible. The Wilker inequality (1.1) has attracted much interest from many mathematicians and has motivated a large number of research papers

involving different proofs, various generalizations and improvements (see [3–5, 8, 11, 14–18, 21, 22, 25–29, 31, 33–35] and the references cited therein).

A related inequality that is of interest to us is Huygens' inequality [12], which asserts that

$$2 \left(\frac{\sin x}{x} \right) + \frac{\tan x}{x} > 3 \quad (0 < |x| < \frac{1}{2}\pi). \quad (1.2)$$

Chen and Cheung [5] showed that for $0 < x < \frac{1}{2}\pi$,

$$3 + \frac{3}{20}x^3 \tan x < 2 \left(\frac{\sin x}{x} \right) + \frac{\tan x}{x} < 3 + \left(\frac{2}{\pi} \right)^4 x^3 \tan x, \quad (1.3)$$

where the constants $\frac{3}{20}$ and $(2/\pi)^4$ are best possible, and obtained a further improvement with the addition of terms involving $x^5 \tan x$. These authors also posed three conjectures on Wilker and Huygens-type inequalities. As far as we know, these conjectures have not yet been proved.

Zhu [32] established some new inequalities of the Huygens-type for trigonometric and hyperbolic functions. Baricz and Sándor [3] pointed out that inequalities (1.1) and (1.2) are simple consequences of the arithmetic-geometric mean inequality, together with the well-known Lazarević-type inequality [13, p. 238] given by $(\cos x)^{1/3} < (\sin x)/x$ ($0 < |x| < \frac{1}{2}\pi$), or equivalently

$$\left(\frac{\sin x}{x} \right)^2 \frac{\tan x}{x} > 1 \quad (0 < |x| < \frac{1}{2}\pi). \quad (1.4)$$

Wu and Srivastava [27, Lemma 3] established another inequality, in which the trigonometric ratios in (1.1) are inverted, namely

$$\left(\frac{x}{\sin x} \right)^2 + \frac{x}{\tan x} > 2 \quad (0 < |x| < \frac{1}{2}\pi) \quad (1.5)$$

and Chen and Sándor [8] established the following inequality chain:

$$\begin{aligned} \frac{(\sin x/x)^2 + \tan x/x}{2} &> \left(\frac{\sin x}{x} \right)^2 \left(\frac{\tan x}{x} \right) > \frac{2(\sin x/x) + \tan x/x}{3} \\ &> \left(\frac{\sin x}{x} \right)^{2/3} \left(\frac{\tan x}{x} \right)^{1/3} > \frac{1}{2} \left[\left(\frac{x}{\sin x} \right)^2 + \frac{x}{\tan x} \right] > \frac{2(x/\sin x) + x/\tan x}{3} > 1 \end{aligned} \quad (1.6)$$

for $0 < |x| < \frac{1}{2}\pi$. This can be seen to involve the Wilker and Huygens inequalities (1.1) and (1.2), together with their inverted forms and the inequality (1.4).

The final inequality we consider is the Papenfuss-Bach inequality. In [20], Papenfuss proposed the following problem to establish the inequality

$$x \sec^2 x - \tan x \leq \frac{8\pi^2 x^3}{(\pi^2 - 4x^2)^2}, \quad 0 \leq x < \frac{1}{2}\pi. \quad (1.7)$$

Bach [2] proved (1.7) and obtained a sharper upper bound in which the numerator $8\pi^2 x^3$ is replaced by $2\pi^4/3$. Ge [9, Theorem 1.3] presented a lower bound in (1.7) and proved that

$$\frac{64x^3}{(\pi^2 - 4x^2)^2} < x \sec^2 x - \tan x \leq \frac{(2\pi^4/3)x^3}{(\pi^2 - 4x^2)^2}, \quad 0 \leq x < \frac{1}{2}\pi, \quad (1.8)$$

where the constants 64 and $2\pi^4/3$ are best possible.

Sun and Zhu [23, Theorem 1.5] obtained better lower and upper bounds for the Papenfuss-Bach inequality in the form

$$\frac{\frac{2\pi^4}{3}x^3 + \left(\frac{8\pi^4}{15} - \frac{16\pi^2}{3} \right) x^5}{(\pi^2 - 4x^2)^2} < x \sec^2 x - \tan x < \frac{\frac{2\pi^4}{3}x^3 + \left(\frac{256}{\pi^2} \alpha - \frac{8\pi^2}{3} \right) x^5}{(\pi^2 - 4x^2)^2} \quad (1.9)$$

with $\alpha = 513/511$. They also posed the following open problem:

Open problem 1. Let $0 < x < \frac{1}{2}\pi$. Then (1.9) holds with $\alpha = 1$, where the constants $\frac{8\pi^4}{15} - \frac{16\pi^2}{3}$ and $\frac{256}{\pi^2} - \frac{8\pi^2}{3}$ are best possible.

Our aim in Sections 2 and 3 is to develop the inverted forms of the Wilker and Huygens inequalities given in (1.5) and the last inequality in (1.6) to produce sharp inequalities. In Section 4, we present a series representation of the remainder in the expansion for $t \sec^2 t - \tan t$. Based on this representation, we establish new bounds for $x \sec^2 x - \tan x$ when $0 < x < \frac{1}{2}\pi$. We also answer the open problem in (1.9).

2. A Wilker-type inequality

We first consider the Wilker-type inequality in (1.5). It is well known [10, p. 42] that

$$\cot x = \frac{1}{x} - \sum_{k=1}^{\infty} \frac{2^{2k} |B_{2k}|}{(2k)!} x^{2k-1}, \quad |x| < \pi. \quad (2.1)$$

Differentiating the expression in (2.1), we find

$$\left(\frac{x}{\sin x}\right)^2 = 1 + \sum_{k=1}^{\infty} \frac{2^{2k}(2k-1)|B_{2k}|}{(2k)!} x^{2k}, \quad |x| < \pi. \quad (2.2)$$

From (2.1) and (2.2), we obtain that for $|x| < \pi$,

$$\left(\frac{x}{\sin x}\right)^2 + \frac{x}{\tan x} = 2 + \sum_{k=1}^{\infty} \frac{k \cdot 2^{2k+3} |B_{2k+2}|}{(2k+2)!} x^{2k+2}. \quad (2.3)$$

It follows from (2.3) that for every $N \in \mathbf{N}$,

$$\frac{N \cdot 2^{2N+3} |B_{2N+2}|}{(2N+2)!} x^{2N+2} < \left(\frac{x}{\sin x}\right)^2 + \frac{x}{\tan x} - \left(2 + \sum_{k=1}^{N-1} \frac{k \cdot 2^{2k+3} |B_{2k+2}|}{(2k+2)!} x^{2k+2}\right) \quad (2.4)$$

for $0 < |x| < \pi$.

In view of (2.4) it is natural to ask: What are the largest number λ_N and the smallest number μ_N such that the inequality

$$\lambda_N x^{2N+2} < \left(\frac{x}{\sin x}\right)^2 + \frac{x}{\tan x} - \left(2 + \sum_{k=1}^{N-1} \frac{k \cdot 2^{2k+3} |B_{2k+2}|}{(2k+2)!} x^{2k+2}\right) < \mu_N x^{2N+2}$$

holds for $x \in (0, \frac{1}{2}\pi)$ and $N \in \mathbf{N}$? Theorem 1 answers this question. In what follows we shall require the summations

$$\begin{aligned} \sum_{k=1}^{\infty} \frac{1}{k^{2n}} &= \frac{2^{2n-1} \pi^{2n}}{(2n)!} |B_{2n}|, & \sum_{k=1}^{\infty} \frac{(-1)^{k+1}}{k^{2n}} &= \frac{(2^{2n-1} - 1) \pi^{2n}}{(2n)!} |B_{2n}|, \\ \sum_{k=1}^{\infty} \frac{1}{(2k-1)^{2n}} &= \frac{(2^n - 1) \pi^{2n}}{2(2n)!} |B_{2n}|, \end{aligned} \quad (2.5)$$

where B_n denote the Bernoulli numbers.

Theorem 1. Let $N \geq 1$ be an integer. Then for $0 < t < \frac{1}{2}\pi$,

$$\lambda_N t^{2N+2} < \left(\frac{t}{\sin t}\right)^2 + \frac{t}{\tan t} - \left(2 + \sum_{j=1}^{N-1} \frac{j \cdot 2^{2j+3} |B_{2j+2}|}{(2j+2)!} t^{2j+2}\right) < \mu_N t^{2N+2} \quad (2.6)$$

with the best possible constants

$$\lambda_N = \frac{N \cdot 2^{2N+3} |B_{2N+2}|}{(2N+2)!} \quad (2.7)$$

and

$$\mu_N = \frac{64N}{\pi^{2N+2}} \sum_{k=1}^{\infty} \frac{1}{k^{2N-2}(4k^2-1)^2} - \frac{16(N-1)}{\pi^{2N+2}} \sum_{k=1}^{\infty} \frac{1}{k^{2N}(4k^2-1)^2}. \quad (2.8)$$

Proof. It follows from [19, p. 118] that

$$\csc^2 t = \frac{1}{t^2} + 2 \sum_{k=1}^{\infty} \frac{t^2 + \pi^2 k^2}{(t^2 - \pi^2 k^2)^2}, \quad \cot t = \frac{1}{t} + 2t \sum_{k=1}^{\infty} \frac{1}{t^2 - \pi^2 k^2}. \quad (2.9)$$

This then yields

$$\left(\frac{t}{\sin t} \right)^2 = 1 + \sum_{k=1}^{\infty} \frac{2t^4 + 2\pi^2 k^2 t^2}{(t^2 - \pi^2 k^2)^2}. \quad (2.10)$$

From (2.10) and the second expansion in (2.9), we obtain

$$\left(\frac{t}{\sin t} \right)^2 + \frac{t}{\tan t} = 2 + 4t^4 \sum_{k=1}^{\infty} \frac{1}{(\pi^2 k^2 - t^2)^2} = 2 + 4t^4 \sum_{k=1}^{\infty} \frac{1}{\pi^4 k^4 (1 - (\frac{t}{\pi k})^2)^2}. \quad (2.11)$$

Using the following identity:

$$\frac{1}{(1-q)^2} = \sum_{j=1}^{N-1} jq^{j-1} + \frac{Nq^{N-1}}{1-q} + \frac{q^N}{(1-q)^2} \quad (q \neq 1) \quad (2.12)$$

and the first sum in (2.5), we then have

$$\begin{aligned} \left(\frac{t}{\sin t} \right)^2 + \frac{t}{\tan t} &= 2 + 4t^4 \sum_{k=1}^{\infty} \frac{1}{\pi^4 k^4 (1 - (\frac{t}{\pi k})^2)^2} \\ &= 2 + 4t^4 \sum_{k=1}^{\infty} \frac{1}{\pi^4 k^4} \left(\sum_{j=1}^{N-1} j \left(\frac{t}{\pi k} \right)^{2j-2} + \frac{N(\frac{t}{\pi k})^{2N-2}}{1 - (\frac{t}{\pi k})^2} + \frac{(\frac{t}{\pi k})^{2N}}{(1 - (\frac{t}{\pi k})^2)^2} \right) \\ &= 2 + \sum_{j=1}^{N-1} \frac{j \cdot 2^{2j+3} |B_{2j+2}|}{(2j+2)!} t^{2j+2} + \sum_{k=1}^{\infty} \frac{4Nt^{2N+2}}{\pi^{2N} k^{2N} (\pi^2 k^2 - t^2)} + \sum_{k=1}^{\infty} \frac{4t^{2N+4}}{\pi^{2N} k^{2N} (\pi^2 k^2 - t^2)^2} \\ &= 2 + \sum_{j=1}^{N-1} \frac{j \cdot 2^{2j+3} |B_{2j+2}|}{(2j+2)!} t^{2j+2} + \frac{4t^{2N+2}}{\pi^{2N}} V_N(t), \end{aligned}$$

where

$$V_N(t) = \sum_{k=1}^{\infty} \frac{N\pi^2 k^2 - (N-1)t^2}{k^{2N} (\pi^2 k^2 - t^2)^2}.$$

Differentiation yields

$$V'_N(t) = \sum_{k=1}^{\infty} \frac{2t \left((N+1)\pi^2 k^2 - (N-1)t^2 \right)}{k^{2N} (\pi^2 k^2 - t^2)^3} > 0.$$

Hence, $V_N(t)$ is strictly increasing for $t \in (0, \frac{1}{2}\pi)$, and we have

$$\lambda_N t^{2N+2} < \left(\frac{t}{\sin t} \right)^2 + \frac{t}{\tan t} - \left(2 + \sum_{j=1}^{N-1} \frac{j \cdot 2^{2j+3} |B_{2j+2}|}{(2j+2)!} t^{2j+2} \right) < \mu_N t^{2N+2}$$

with

$$\lambda_N = \frac{4}{\pi^{2N}} V_N(0) \quad \text{and} \quad \mu_N = \frac{4}{\pi^{2N}} V_N\left(\frac{1}{2}\pi\right).$$

Direct computations yield

$$V_N(0) = \frac{N}{\pi^2} \sum_{k=1}^{\infty} \frac{1}{k^{2N+2}} = \frac{N \cdot 2^{2N+1} \pi^{2N} |B_{2N+2}|}{(2N+2)!}$$

and

$$V_N\left(\frac{1}{2}\pi\right) = \frac{16N}{\pi^2} \sum_{k=1}^{\infty} \frac{1}{k^{2N-2}(4k^2-1)^2} - \frac{4(N-1)}{\pi^2} \sum_{k=1}^{\infty} \frac{1}{k^{2N}(4k^2-1)^2}.$$

Hence, the inequality (2.6) holds with the best possible constants given in (2.7) and (2.8). The proof of Theorem 1 is complete. \square

Remark 1. Direct computations yield

$$\lambda_1 = \frac{2}{45}, \quad \mu_1 = \frac{4(\pi^2 - 8)}{\pi^4}$$

and

$$\lambda_2 = \frac{8}{945}, \quad \mu_2 = \frac{8(-720 + 90\pi^2 - \pi^4)}{45\pi^6}.$$

We then obtain from (2.6) that for $0 < t < \frac{1}{2}\pi$,

$$2 + \frac{2}{45}t^4 < \left(\frac{t}{\sin t}\right)^2 + \frac{t}{\tan t} < 2 + \frac{4(\pi^2 - 8)}{\pi^4}t^4, \quad (2.13)$$

where the constants $\frac{2}{45}$ and $4(\pi^2 - 8)/\pi^4$ are best possible, and

$$2 + \frac{2}{45}t^4 + \frac{8}{945}t^6 < \left(\frac{t}{\sin t}\right)^2 + \frac{t}{\tan t} < 2 + \frac{2}{45}t^4 + \frac{8(-720 + 90\pi^2 - \pi^4)}{45\pi^6}t^6, \quad (2.14)$$

where the constants $\frac{8}{945}$ and $8(-720 + 90\pi^2 - \pi^4)/(45\pi^6)$ are best possible.

The classical Euler gamma function is defined (for $x > 0$) by

$$\Gamma(x) = \int_0^{\infty} t^{x-1} e^{-t} dt.$$

Its logarithmic derivative, denoted by $\psi(x) = \Gamma'(x)/\Gamma(x)$, is called the psi (or digamma) function, and $\psi^{(k)}(x)$ ($k \in \mathbf{N}$) are called the polygamma functions.

Theorem 2. Let $N \geq 0$ be an integer. Then for $0 < x < \frac{1}{2}\pi$,

$$\alpha_N x^4 < \left(\frac{x}{\sin x}\right)^2 + \frac{x}{\tan x} - \left(2 + 4x^4 \sum_{k=1}^N \frac{1}{(\pi^2 k^2 - x^2)^2}\right) < \beta_N x^4 \quad (2.15)$$

with the best possible constants

$$\alpha_N = \frac{2\psi'''(N+1)}{3\pi^4} \quad \text{and} \quad \beta_N = \frac{8\left((2N+1)^2\psi'(N+\frac{1}{2}) - 4(N+1)\right)}{(2N+1)^2\pi^4}. \quad (2.16)$$

Proof. Write (2.11) as

$$\left(\frac{x}{\sin x}\right)^2 + \frac{x}{\tan x} = 2 + 4x^4 \sum_{k=1}^N \frac{1}{(\pi^2 k^2 - x^2)^2} + 4x^4 A_N(x),$$

where

$$A_N(x) = \sum_{k=N+1}^{\infty} \frac{1}{(\pi^2 k^2 - x^2)^2}.$$

Obviously, $A_N(x)$ is strictly increasing for $x \in (0, \frac{1}{2}\pi)$. Hence, for $0 < x < \frac{1}{2}\pi$, we have

$$\alpha_N x^4 < \left(\frac{x}{\sin x}\right)^2 + \frac{x}{\tan x} - \left(2 + 4x^4 \sum_{k=1}^N \frac{1}{(\pi^2 k^2 - x^2)^2}\right) < \beta_N x^4$$

with

$$\alpha_N = 4A_N(0) = \frac{4}{\pi^4} \sum_{k=N+1}^{\infty} \frac{1}{k^4} \quad \text{and} \quad \beta_N = 4A_N\left(\frac{1}{2}\pi\right) = \frac{64}{\pi^4} \sum_{k=N+1}^{\infty} \frac{1}{(4k^2 - 1)^2}.$$

From the following formula (see [1, p. 260, Eq. (6.4.10)]):

$$\psi^{(n)}(z) = (-1)^{n+1} n! \sum_{k=0}^{\infty} \frac{1}{(z+k)^{n+1}}, \quad z \neq 0, -1, -2, \dots,$$

we obtain

$$\sum_{k=N+1}^{\infty} \frac{1}{k^4} = \frac{\psi'''(N+1)}{6}. \quad (2.17)$$

We find¹

$$\sum_{k=N+1}^{\infty} \frac{1}{(4k^2 - 1)^2} = \frac{1}{8} \psi' \left(N + \frac{1}{2} \right) - \frac{N+1}{2(2N+1)^2}. \quad (2.18)$$

Hence, the inequality (2.15) holds with the best possible constants given in (2.16). The proof of Theorem 2 is complete. \square

Remark 2. The choice $N = 0$ in (2.15) yields (2.13). The choice $N = 1$ in (2.15) yields

$$2 + \frac{4x^4}{(\pi^2 - x^2)^2} + \frac{2(\pi^4 - 90)}{45\pi^4} x^4 < \left(\frac{x}{\sin x}\right)^2 + \frac{x}{\tan x} < 2 + \frac{4x^4}{(\pi^2 - x^2)^2} + \frac{4(9\pi^2 - 88)}{9\pi^4} x^4 \quad (2.19)$$

for $0 < x < \frac{1}{2}\pi$, where the constants $2(\pi^4 - 90)/(45\pi^4)$ and $4(9\pi^2 - 88)/(9\pi^4)$ are best possible.

Remark 3. There is no strict comparison between the two lower bounds in (2.14) and (2.19). Likewise, there is no strict comparison between the two upper bounds in (2.14) and (2.19).

Theorem 3 proves Conjecture 2 in [5].

Theorem 3. Let $N \geq 1$ be an integer. Then for $0 < x < \frac{1}{2}\pi$, we have

$$\begin{aligned} 2 + \sum_{k=1}^{N-1} \frac{k \cdot 2^{2k+3} |B_{2k+2}|}{(2k+2)!} x^{2k+2} + p_N x^{2N+1} \tan x &< \left(\frac{x}{\sin x}\right)^2 + \frac{x}{\tan x} \\ &< 2 + \sum_{k=1}^{N-1} \frac{k \cdot 2^{2k+3} |B_{2k+2}|}{(2k+2)!} x^{2k+2} + q_N x^{2N+1} \tan x \end{aligned} \quad (2.20)$$

with the best possible constants

$$p_N = 0 \quad \text{and} \quad q_N = \frac{N \cdot 2^{2N+3} |B_{2N+2}|}{(2N+2)!}. \quad (2.21)$$

¹The formula (2.18) is established by induction on N in the appendix.

Proof. By (2.3), for $p_N = 0$, the first inequality in (2.20) holds. We now prove the second inequality in (2.20) with $q_N = N \cdot 2^{2N+3} |B_{2N+2}| / (2N+2)!$. Using (2.3) and the following expansion (see [10, p. 42]):

$$\tan x = \sum_{k=1}^{\infty} \frac{2^{2k}(2^{2k}-1)|B_{2k}|}{(2k)!} x^{2k-1}, \quad |x| < \frac{1}{2}\pi, \quad (2.22)$$

we find

$$\begin{aligned} & \frac{N \cdot 2^{2N+3} |B_{2N+2}|}{(2N+2)!} x^{2N+1} \tan x - \left(\left(\frac{x}{\sin x} \right)^2 + \frac{x}{\tan x} - 2 - \sum_{k=1}^{N-1} \frac{k \cdot 2^{2k+3} |B_{2k+2}|}{(2k+2)!} x^{2k+2} \right) \\ &= \frac{N \cdot 2^{2N+3} |B_{2N+2}|}{(2N+2)!} x^{2N+1} \sum_{k=1}^{\infty} \frac{2^{2k}(2^{2k}-1)|B_{2k}|}{(2k)!} x^{2k-1} - \sum_{k=N+1}^{\infty} \frac{(k-1) \cdot 2^{2k+1} |B_{2k}|}{(2k)!} x^{2k} \\ &= \sum_{k=N+2}^{\infty} \left\{ \frac{N \cdot 2^{2N+3} |B_{2N+2}|}{(2N+2)!} \frac{2^{2k-2N}(2^{2k-2N}-1)|B_{2k-2N}|}{(2k-2N)!} - \frac{(k-1) \cdot 2^{2k+1} |B_{2k}|}{(2k)!} \right\} x^{2k}, \quad (2.23) \end{aligned}$$

where we note that the term corresponding to $k = N+1$ vanishes.

We claim that for $k \geq N+2$,

$$\frac{N \cdot 2^{2N+3} |B_{2N+2}|}{(2N+2)!} \frac{2^{2k-2N}(2^{2k-2N}-1)|B_{2k-2N}|}{(2k-2N)!} > \frac{(k-1) \cdot 2^{2k+1} |B_{2k}|}{(2k)!}. \quad (2.24)$$

Using the inequality (see [1, p. 805])

$$\frac{2}{(2\pi)^{2n} (1-2^{1-2n})} > \frac{|B_{2n}|}{(2n)!} > \frac{2}{(2\pi)^{2n}}, \quad n \geq 1, \quad (2.25)$$

it is sufficient to prove that for $k \geq N+2$,

$$\frac{N \cdot 2^{2N+3} \cdot 2 \cdot 2^{2k-2N} (2^{2k-2N}-1) \cdot 2}{(2\pi)^{2N+2} (2\pi)^{2k-2N}} > \frac{2(k-1) \cdot 2^{2k+1}}{(2\pi)^{2k} (1-2^{1-2k})},$$

which can be rearranged as

$$N \left(\frac{2^{2k}}{2^{2N}} - 1 \right) > \frac{\pi^2}{2} (k-1) \left(1 + \frac{2}{2^{2k}-2} \right), \quad k \geq N+2.$$

Noting that $\pi^2/2 < 5$, it is enough to prove the following inequality:

$$N \left(\frac{2^{2k}}{2^{2N}} - 1 \right) > 5(k-1) \left(1 + \frac{2}{2^{2k}-2} \right), \quad k \geq N+2,$$

which can be rearranged as

$$\frac{N}{2^{2N}} 2^{2k} - 5(k-1) > N + \frac{10(k-1)}{2^{2k}-2}, \quad k \geq N+2.$$

Noting that the sequence

$$\frac{N}{2^{2N}} 2^{2k} - 5(k-1)$$

is strictly increasing for $k \geq N+2$, and the sequence

$$\frac{10(k-1)}{2^{2k}-2}$$

is strictly decreasing for $k \geq 2$, it is enough to prove the following inequality:

$$\frac{N}{2^{2N}} 2^{2(N+2)} - 5(N+1) > N + \frac{10(N+1)}{2^{2(N+2)}-2},$$

which can be rearranged as

$$(2N - 1)2^{2N+3} > 3N, \quad N \geq 1.$$

Obviously, the last inequality holds. This proves the claim (2.24). From (2.23), we obtain the second inequality in (2.20) with $q_N = N \cdot 2^{2N+3}|B_{2N+2}|/(2N+2)!$.

Write (2.20) as

$$p_N < \frac{\left(\frac{x}{\sin x}\right)^2 + \frac{x}{\tan x} - 2 - \sum_{k=1}^{N-1} \frac{k \cdot 2^{2k+3} |B_{2k+2}|}{(2k+2)!} x^{2k+2}}{x^{2N+1} \tan x} < q_N.$$

We then find that

$$\lim_{x \rightarrow \frac{\pi}{2}} \frac{\left(\frac{x}{\sin x}\right)^2 + \frac{x}{\tan x} - 2 - \sum_{k=1}^{N-1} \frac{k \cdot 2^{2k+3} |B_{2k+2}|}{(2k+2)!} x^{2k+2}}{x^{2N+1} \tan x} = 0$$

and

$$\lim_{x \rightarrow 0} \frac{\left(\frac{x}{\sin x}\right)^2 + \frac{x}{\tan x} - 2 - \sum_{k=1}^{N-1} \frac{k \cdot 2^{2k+3} |B_{2k+2}|}{(2k+2)!} x^{2k+2}}{x^{2N+1} \tan x} = \frac{N \cdot 2^{2N+3} |B_{2N+2}|}{(2N+2)!}.$$

Hence, the inequality (2.20) holds with the best possible constants given in (2.21). The proof of Theorem 3 is complete. \square

3. A Huygens-type inequality

We now turn our attention to the inverted form of the Huygens inequality in (1.2). Using (2.1) and the following expansion (see [10, p. 43]):

$$\csc x = \frac{1}{x} + \sum_{k=1}^{\infty} \frac{2(2^{2k-1} - 1)|B_{2k}|}{(2k)!} x^{2k-1}, \quad |x| < \pi,$$

we find

$$2\left(\frac{x}{\sin x}\right) + \frac{x}{\tan x} = 3 + \sum_{k=2}^{\infty} \frac{(2^{2k} - 4)|B_{2k}|}{(2k)!} x^{2k}, \quad |x| < \pi. \quad (3.1)$$

It follows from (3.1) that for every $N \in \mathbf{N}$,

$$\frac{(2^{2N+2} - 4)|B_{2N+2}|}{(2N+2)!} x^{2N+2} < 2\left(\frac{x}{\sin x}\right) + \frac{x}{\tan x} - \left(3 + \sum_{k=2}^N \frac{(2^{2k} - 4)|B_{2k}|}{(2k)!} x^{2k}\right) \quad (3.2)$$

for $0 < |x| < \pi$.

In view of (3.2) it is natural to ask: What are the largest number a_N and the smallest number b_N such that the inequality

$$a_N x^{2N+2} < 2\left(\frac{x}{\sin x}\right) + \frac{x}{\tan x} - \left(3 + \sum_{k=2}^N \frac{(2^{2k} - 4)|B_{2k}|}{(2k)!} x^{2k}\right) < b_N x^{2N+2}$$

holds for $x \in (0, \frac{1}{2}\pi)$ and $N \in \mathbf{N}$? Theorem 4 answers this question.

Theorem 4. *Let $N \geq 1$ be an integer. Then for $0 < |x| < \frac{1}{2}\pi$,*

$$a_N x^{2N+2} < 2\left(\frac{x}{\sin x}\right) + \frac{x}{\tan x} - \left(3 + \sum_{j=2}^N \frac{(2^{2j} - 4)|B_{2j}|}{(2j)!} x^{2j}\right) < b_N x^{2N+2} \quad (3.3)$$

with the best possible constants

$$a_N = \frac{(2^{2N+2} - 4)|B_{2N+2}|}{(2N+2)!} \quad (3.4)$$

and

$$b_N = \frac{8}{\pi^{2N+2}} \left(\sum_{k=1}^{\infty} \frac{(-1)^{k+1}}{k^{2N}(2k-1)} - \sum_{k=1}^{\infty} \frac{(-1)^{k+1}}{k^{2N}(2k+1)} \right) - \frac{4}{\pi^{2N+2}} \left(\sum_{k=1}^{\infty} \frac{1}{k^{2N}(2k-1)} - \sum_{k=1}^{\infty} \frac{1}{k^{2N}(2k+1)} \right). \quad (3.5)$$

Proof. By [6, Theorems 4, 5], we have

$$2 \left(\frac{x}{\sin x} \right) = 2 + \sum_{j=1}^N \frac{(2^{2j+1} - 4)|B_{2j}|}{(2j)!} x^{2j} + x^{2N+2} \sum_{k=1}^{\infty} \frac{4(-1)^{k+1}}{(k\pi)^{2N}((k\pi)^2 - x^2)} \quad (3.6)$$

and

$$\frac{x}{\tan x} = 1 - \sum_{j=1}^N \frac{2^{2j}|B_{2j}|}{(2j)!} x^{2j} - x^{2N+2} \sum_{k=1}^{\infty} \frac{2}{(k\pi)^{2N}((k\pi)^2 - x^2)}. \quad (3.7)$$

Adding these two expressions, we obtain

$$2 \left(\frac{x}{\sin x} \right) + \frac{x}{\tan x} = 3 + \sum_{j=2}^N \frac{(2^{2j} - 4)|B_{2j}|}{(2j)!} x^{2j} + \frac{2x^{2N+2}}{\pi^{2N}} U_N(x),$$

where

$$U_N(x) = \sum_{k=1}^{\infty} (-1)^{k+1} \frac{2 - (-1)^{k+1}}{k^{2N}((k\pi)^2 - x^2)}.$$

Differentiation yields

$$\frac{U'_N(x)}{2x} = \sum_{k=1}^{\infty} (-1)^{k+1} \alpha_k, \quad \alpha_k = \frac{2 - (-1)^{k+1}}{k^{2N}((k\pi)^2 - x^2)^2}. \quad (3.8)$$

Then it is easily seen that $\alpha_k > \alpha_{k+1}$ for $k \in \mathbf{N}$, $0 < x < \frac{1}{2}\pi$ and $N \in \mathbf{N}$; thus for every $N \geq 1$, we have $U'_N(x) > 0$ for $0 < x < \frac{1}{2}\pi$. Hence, for all $0 < x < \frac{1}{2}\pi$ and $N \in \mathbf{N}$, we have

$$U_N(0) < U_N(x) < U_N\left(\frac{1}{2}\pi\right).$$

Using (2.5), we find

$$\begin{aligned} a_N &= \frac{2U_N(0)}{\pi^{2N}} = 4 \sum_{k=1}^{\infty} \frac{(-1)^{k+1}}{(k\pi)^{2N+2}} - 2 \sum_{k=1}^{\infty} \frac{1}{(k\pi)^{2N+2}} \\ &= \frac{4(2^{2N+1} - 1)}{(2N+2)!} |B_{2N+2}| - \frac{2 \cdot 2^{2N+1}}{(2N+2)!} |B_{2N+2}| = \frac{(2^{2N+2} - 4)|B_{2N+2}|}{(2N+2)!} \end{aligned}$$

and

$$\begin{aligned} b_N &= \frac{2U_N\left(\frac{1}{2}\pi\right)}{\pi^{2N}} = 4 \sum_{k=1}^{\infty} \frac{(-1)^{k+1}}{(k\pi)^{2N}((k\pi)^2 - (\pi/2)^2)} - 2 \sum_{k=1}^{\infty} \frac{1}{(k\pi)^{2N}((k\pi)^2 - (\pi/2)^2)} \\ &= \frac{8}{\pi^{2N+2}} \left(\sum_{k=1}^{\infty} \frac{(-1)^{k+1}}{k^{2N}(2k-1)} - \sum_{k=1}^{\infty} \frac{(-1)^{k+1}}{k^{2N}(2k+1)} \right) \\ &\quad - \frac{4}{\pi^{2N+2}} \left(\sum_{k=1}^{\infty} \frac{1}{k^{2N}(2k-1)} - \sum_{k=1}^{\infty} \frac{1}{k^{2N}(2k+1)} \right). \end{aligned}$$

The proof of Theorem 4 is complete. \square

Clearly,

$$a_1 = \frac{1}{60} \quad \text{and} \quad a_2 = \frac{1}{504}.$$

Direct computations yield

$$\begin{aligned} \sum_{k=1}^{\infty} \frac{(-1)^{k+1}}{k^2(2k-1)} &= \pi - 2 \ln 2 - \frac{\pi^2}{12}, & \sum_{k=1}^{\infty} \frac{(-1)^{k+1}}{k^2(2k+1)} &= 4 - 2 \ln 2 - \pi + \frac{\pi^2}{12}, \\ \sum_{k=1}^{\infty} \frac{1}{k^2(2k-1)} &= -\frac{\pi^2}{6} + 4 \ln 2, & \sum_{k=1}^{\infty} \frac{1}{k^2(2k+1)} &= -4 + 4 \ln 2 + \frac{\pi^2}{6}, \\ \sum_{k=1}^{\infty} \frac{(-1)^{k+1}}{k^4(2k-1)} &= 4\pi - 8 \ln 2 - \frac{\pi^2}{3} - \frac{3}{2}\zeta(3) - \frac{7\pi^4}{720}, \\ \sum_{k=1}^{\infty} \frac{(-1)^{k+1}}{k^4(2k+1)} &= 16 - 4\pi - 8 \ln 2 + \frac{\pi^2}{3} - \frac{3}{2}\zeta(3) + \frac{7\pi^4}{720}, \\ \sum_{k=1}^{\infty} \frac{1}{k^4(2k-1)} &= 16 \ln 2 - \frac{2\pi^2}{3} - 2\zeta(3) - \frac{\pi^4}{90}, \\ \sum_{k=1}^{\infty} \frac{1}{k^4(2k+1)} &= -16 + 16 \ln 2 + \frac{2\pi^2}{3} - 2\zeta(3) + \frac{\pi^4}{90}, \end{aligned}$$

where $\zeta(s)$ is the Riemann zeta function. Then, we obtain from (3.5)

$$b_1 = \frac{16(\pi-3)}{\pi^4} \quad \text{and} \quad b_2 = \frac{960\pi - \pi^4 - 2880}{15\pi^6}.$$

From (3.3), we have, for $0 < |x| < \frac{1}{2}\pi$,

$$3 + \frac{1}{60}x^4 < 2 \left(\frac{x}{\sin x} \right) + \frac{x}{\tan x} < 3 + \frac{16(\pi-3)}{\pi^4}x^4, \quad (3.9)$$

where the constants $\frac{1}{60}$ and $16(\pi-3)/\pi^4$ are best possible, and

$$3 + \frac{1}{60}x^4 + \frac{1}{504}x^6 < 2 \left(\frac{x}{\sin x} \right) + \frac{x}{\tan x} < 3 + \frac{1}{60}x^4 + \frac{960\pi - \pi^4 - 2880}{15\pi^6}x^6, \quad (3.10)$$

where the constants $\frac{1}{504}$ and $(960\pi - \pi^4 - 2880)/(15\pi^6)$ are best possible.

The formula (3.1) motivated us to observe Theorem 5.

Theorem 5. *Let $N \geq 1$ be an integer. Then for $0 < x < \frac{1}{2}\pi$, we have*

$$\begin{aligned} 3 + \sum_{j=2}^N \frac{(2^{2j}-4)|B_{2j}|}{(2j)!} x^{2j} + \rho_N x^{2N+1} \tan x &< 2 \left(\frac{x}{\sin x} \right) + \frac{x}{\tan x} \\ &< 3 + \sum_{j=2}^N \frac{(2^{2j}-4)|B_{2j}|}{(2j)!} x^{2j} + \varrho_N x^{2N+1} \tan x \end{aligned} \quad (3.11)$$

with the best possible constants

$$\rho_N = 0 \quad \text{and} \quad \varrho_N = \frac{4(2^{2N}-1)|B_{2N+2}|}{(2N+2)!}. \quad (3.12)$$

Proof. By (3.1), for $\rho_N = 0$, the first inequality in (3.11) holds. We now prove the second inequality in (3.12) with $\varrho_N = 4(2^{2N} - 1)|B_{2N+2}|/(2N + 2)!$. Using the expansion [10, p. 44]

$$\tan t = \sum_{k=1}^{\infty} \frac{8t}{\pi^2(2k-1)^2 - 4t^2} \quad (3.13)$$

and (3.1), we find

$$\begin{aligned} & \frac{4(2^{2N} - 1)|B_{2N+2}|}{(2N + 2)!} x^{2N+1} \tan x - \left(2 \left(\frac{x}{\sin x} \right) + \frac{x}{\tan x} - 3 - \sum_{j=2}^N \frac{(2^{2j} - 4)|B_{2j}|}{(2j)!} x^{2j} \right) \\ &= \frac{4(2^{2N} - 1)|B_{2N+2}|}{(2N + 2)!} x^{2N+1} \sum_{k=1}^{\infty} \frac{2^{2k}(2^{2k} - 1)|B_{2k}|}{(2k)!} x^{2k-1} - \sum_{k=N+1}^{\infty} \frac{(2^{2k} - 4)|B_{2k}|}{(2k)!} x^{2k} \\ &= \sum_{k=N+2}^{\infty} \left\{ \frac{4(2^{2N} - 1)|B_{2N+2}|}{(2N + 2)!} \frac{2^{2k-2N}(2^{2k-2N} - 1)|B_{2k-2N}|}{(2k - 2N)!} - \frac{(2^{2k} - 4)|B_{2k}|}{(2k)!} \right\} x^{2k}. \end{aligned} \quad (3.14)$$

We claim that for $k \geq N + 2$,

$$\frac{4(2^{2N} - 1)|B_{2N+2}|}{(2N + 2)!} \frac{2^{2k-2N}(2^{2k-2N} - 1)|B_{2k-2N}|}{(2k - 2N)!} > \frac{(2^{2k} - 4)|B_{2k}|}{(2k)!}. \quad (3.15)$$

Using the inequality (2.25), it is sufficient to prove that

$$\frac{4(2^{2N} - 1) \cdot 2 \cdot 2^{2k-2N}(2^{2k-2N} - 1) \cdot 2}{(2\pi)^{2N+2} (2\pi)^{2k-2N}} > \frac{(2^{2k} - 4) \cdot 2}{(2\pi)^{2k} (1 - 2^{1-2k})}, \quad k \geq N + 2,$$

which can be rearranged as

$$\left(1 - \frac{1}{2^{2N}} \right) \left(\frac{2^{2k}}{2^{2N}} - 1 \right) > \frac{\pi^2}{2} \left(1 - \frac{2}{2^{2k} - 2} \right), \quad k \geq N + 2.$$

Noting that $\pi^2/2 < 5$, it is enough to prove the following inequality:

$$\left(1 - \frac{1}{2^{2N}} \right) \left(\frac{2^{2k}}{2^{2N}} - 1 \right) > 5 \left(1 - \frac{2}{2^{2k} - 2} \right), \quad k \geq N + 2,$$

which can be written as

$$\left(1 - \frac{1}{2^{2N}} \right) \frac{2^{2k}}{2^{2N}} + \frac{1}{2^{2N}} + \frac{10}{2^{2k} - 2} > 6, \quad k \geq N + 2.$$

It is enough to prove the following inequality:

$$\left(1 - \frac{1}{2^{2N}} \right) \frac{2^{2k}}{2^{2N}} + \frac{1}{2^{2N}} > 6, \quad k \geq N + 2. \quad (3.16)$$

Clearly,

$$\left(1 - \frac{1}{2^{2N}} \right) \frac{2^{2k}}{2^{2N}} + \frac{1}{2^{2N}} \geq \left(1 - \frac{1}{2^{2N}} \right) \frac{2^{2N+4}}{2^{2N}} + \frac{1}{2^{2N}} = 16 - \frac{15}{2^{2N}}, \quad k \geq N + 2.$$

In order to prove (3.16), it suffices to show that

$$16 - \frac{15}{2^{2N}} > 6, \quad N \geq 1,$$

that is,

$$2^{2N+1} > 3, \quad N \geq 1.$$

Obviously, the last inequality holds. This proves the claim (3.15). From (3.14), we obtain the second inequality in (3.11) with $\varrho_N = 4(2^{2N} - 1)|B_{2N+2}|/(2N + 2)!$.

Write (3.11) as

$$\rho_N < \frac{2\left(\frac{x}{\sin x}\right) + \frac{x}{\tan x} - 3 - \sum_{j=2}^N \frac{(2^{2j}-4)|B_{2j}|}{(2j)!} x^{2j}}{x^{2N+1} \tan x} < \varrho_N.$$

We then find that

$$\lim_{x \rightarrow \frac{\pi}{2}} \frac{2\left(\frac{x}{\sin x}\right) + \frac{x}{\tan x} - 3 - \sum_{j=2}^N \frac{(2^{2j}-4)|B_{2j}|}{(2j)!} x^{2j}}{x^{2N+1} \tan x} = 0$$

and

$$\lim_{x \rightarrow 0} \frac{2\left(\frac{x}{\sin x}\right) + \frac{x}{\tan x} - 3 - \sum_{j=2}^N \frac{(2^{2j}-4)|B_{2j}|}{(2j)!} x^{2j}}{x^{2N+1} \tan x} = \frac{4(2^{2N} - 1)|B_{2N+2}|}{(2N + 2)!}.$$

Hence, the inequality (3.11) holds with the best possible constants given in (3.12). The proof of Theorem 5 is complete. \square

Remark 4. For $0 < |x| < \frac{1}{2}\pi$, we have

$$3 + ax^3 \tan x < 2\left(\frac{x}{\sin x}\right) + \frac{x}{\tan x} < 3 + bx^3 \tan x \quad (3.17)$$

with the best possible constants

$$a = 0 \quad \text{and} \quad b = \frac{1}{60}. \quad (3.18)$$

There is no strict comparison between the two upper bounds in (3.9) and (3.17).

4. The Papenfuss-Bach inequality

In this section we consider refinements to the Papenfuss-Bach inequality stated in (1.7) and answer the open problem in (1.9).

It follows from [10, p. 44] that

$$\sec^2 \frac{\pi x}{2} = \frac{4}{\pi^2} \sum_{k=1}^{\infty} \left\{ \frac{1}{(2k-1-x)^2} + \frac{1}{(2k-1+x)^2} \right\}.$$

Replacement of x by $2t/\pi$ yields

$$\sec^2 t = \frac{4}{\pi^2} \sum_{k=1}^{\infty} \left\{ \frac{1}{(2k-1-\frac{2t}{\pi})^2} + \frac{1}{(2k-1+\frac{2t}{\pi})^2} \right\}. \quad (4.1)$$

From (3.13) and (4.1), we have

$$t \sec^2 t - \tan t = \frac{64t^3}{\pi^4} \sum_{k=1}^{\infty} \frac{1}{(2k-1)^4 \left(1 - \left(\frac{2t}{\pi(2k-1)}\right)^2\right)^2}. \quad (4.2)$$

Using (2.12) and the third summation in (2.5), we obtain from (4.2) the series representation of the remainder in the expansion for $\sec^2 t - \tan t/t$:

$$\begin{aligned} & t \sec^2 t - \tan t \\ &= \frac{64t^3}{\pi^4} \sum_{k=1}^{\infty} \frac{1}{(2k-1)^4} \left(\sum_{j=1}^{N-1} j \left(\frac{2t}{\pi(2k-1)}\right)^{2j-2} + \frac{N \left(\frac{2t}{\pi(2k-1)}\right)^{2N-2}}{1 - \left(\frac{2t}{\pi(2k-1)}\right)^2} + \frac{\left(\frac{2t}{\pi(2k-1)}\right)^{2N}}{\left(1 - \left(\frac{2t}{\pi(2k-1)}\right)^2\right)^2} \right) \\ &= \sum_{j=1}^{N-1} \frac{2j \cdot 2^{2j+2} (2^{2j+2} - 1) |B_{2j+2}|}{(2j+2)!} t^{2j+1} + \kappa_N(t), \end{aligned} \quad (4.3)$$

where

$$\begin{aligned} \kappa_N(t) &= \frac{N \cdot 2^{2N+4} t^{2N+1}}{\pi^{2N}} \sum_{k=1}^{\infty} \frac{1}{(2k-1)^{2N} (\pi^2(2k-1)^2 - 4t^2)} \\ &\quad + \frac{2^{2N+6} t^{2N+3}}{\pi^{2N}} \sum_{k=1}^{\infty} \frac{1}{(2k-1)^{2N} (\pi^2(2k-1)^2 - 4t^2)^2}. \end{aligned} \quad (4.4)$$

Theorem 6. Let $N \geq 1$ be an integer. Then for $0 < t < \frac{1}{2}\pi$, we have

$$\begin{aligned} L_N(t) &< t \sec^2 t - \tan t - \sum_{j=1}^{N-1} \frac{2j \cdot 2^{2j+2} (2^{2j+2} - 1) |B_{2j+2}| t^{2j+1}}{(2j+2)!} \\ &\quad - \frac{N \cdot 2^{2N+4} t^{2N+1}}{\pi^{2N} (\pi^2 - 4t^2)} - \frac{2^{2N+6} t^{2N+3}}{\pi^{2N} (\pi^2 - 4t^2)^2} < M_N(t), \end{aligned} \quad (4.5)$$

where

$$\begin{aligned} L_N(t) &= \frac{N \cdot 2^{2N+4} t^{2N+1}}{\pi^{2N+2}} \left\{ \frac{(2^{2N+2} - 1) \pi^{2N+2} |B_{2N+2}|}{2 \cdot (2N+2)!} - 1 \right\} \\ &\quad + \frac{2^{2N+6} t^{2N+3}}{\pi^{2N+4}} \left\{ \frac{(2^{2N+4} - 1) \pi^{2N+4} |B_{2N+4}|}{2 \cdot (2N+4)!} - 1 \right\} \end{aligned}$$

and

$$M_N(t) = \frac{N \cdot 2^{2N+2} t^{2N+1}}{\pi^{2N+2}} \sum_{k=2}^{\infty} \frac{1}{(2k-1)^{2N} k(k-1)} + \frac{2^{2N+2} t^{2N+3}}{\pi^{2N+4}} \sum_{k=2}^{\infty} \frac{1}{(2k-1)^{2N} k^2(k-1)^2}.$$

Proof. Write (4.3) as

$$\begin{aligned} t \sec^2 t - \tan t &= \sum_{j=1}^{N-1} \frac{2j \cdot 2^{2j+2} (2^{2j+2} - 1) |B_{2j+2}| t^{2j+1}}{(2j+2)!} \\ &\quad + \frac{N \cdot 2^{2N+4} t^{2N+1}}{\pi^{2N} (\pi^2 - 4t^2)} + \frac{N \cdot 2^{2N+4} t^{2N+1}}{\pi^{2N}} I_N(t) \\ &\quad + \frac{2^{2N+6} t^{2N+3}}{\pi^{2N} (\pi^2 - 4t^2)^2} + \frac{2^{2N+6} t^{2N+3}}{\pi^{2N}} J_N(t), \end{aligned} \quad (4.6)$$

where

$$I_N(t) = \sum_{k=2}^{\infty} \frac{1}{(2k-1)^{2N} (\pi^2(2k-1)^2 - 4t^2)}$$

and

$$J_N(t) = \sum_{k=2}^{\infty} \frac{1}{(2k-1)^{2N} (\pi^2(2k-1)^2 - 4t^2)^2}.$$

Obviously, $I_N(t)$ and $J_N(t)$ are both strictly increasing for $t \in (0, \frac{1}{2}\pi)$. We then obtain from (4.6) that

$$\begin{aligned} &\frac{N \cdot 2^{2N+4} t^{2N+1}}{\pi^{2N}} I_N(0) + \frac{2^{2N+6} t^{2N+3}}{\pi^{2N}} J_N(0) \\ &< t \sec^2 t - \tan t - \sum_{j=1}^{N-1} \frac{2j \cdot 2^{2j+2} (2^{2j+2} - 1) |B_{2j+2}| t^{2j+1}}{(2j+2)!} - \frac{N \cdot 2^{2N+4} t^{2N+1}}{\pi^{2N} (\pi^2 - 4t^2)} - \frac{2^{2N+6} t^{2N+3}}{\pi^{2N} (\pi^2 - 4t^2)^2} \\ &< \frac{N \cdot 2^{2N+4} t^{2N+1}}{\pi^{2N}} I_N\left(\frac{1}{2}\pi\right) + \frac{2^{2N+6} t^{2N+3}}{\pi^{2N}} J_N\left(\frac{1}{2}\pi\right). \end{aligned}$$

Direct computations yield

$$\begin{aligned}
L_N(t) &= \frac{N \cdot 2^{2N+4} t^{2N+1}}{\pi^{2N}} I_N(0) + \frac{2^{2N+6} t^{2N+3}}{\pi^{2N}} J_N(0) \\
&= \frac{N \cdot 2^{2N+4} t^{2N+1}}{\pi^{2N+2}} \sum_{k=2}^{\infty} \frac{1}{(2k-1)^{2N+2}} + \frac{2^{2N+6} t^{2N+3}}{\pi^{2N+4}} \sum_{k=2}^{\infty} \frac{1}{(2k-1)^{2N+4}} \\
&= \frac{N \cdot 2^{2N+4} t^{2N+1}}{\pi^{2N+2}} \left\{ \frac{(2^{2N+2} - 1) \pi^{2N+2} |B_{2N+2}|}{2 \cdot (2N+2)!} - 1 \right\} \\
&\quad + \frac{2^{2N+6} t^{2N+3}}{\pi^{2N+4}} \left\{ \frac{(2^{2N+4} - 1) \pi^{2N+4} |B_{2N+4}|}{2 \cdot (2N+4)!} - 1 \right\}
\end{aligned}$$

and

$$\begin{aligned}
M_N(t) &= \frac{N \cdot 2^{2N+4} t^{2N+1}}{\pi^{2N}} I_N\left(\frac{1}{2}\pi\right) + \frac{2^{2N+6} t^{2N+3}}{\pi^{2N}} J_N\left(\frac{1}{2}\pi\right) \\
&= \frac{N \cdot 2^{2N+2} t^{2N+1}}{\pi^{2N+2}} \sum_{k=2}^{\infty} \frac{1}{(2k-1)^{2N} k(k-1)} + \frac{2^{2N+2} t^{2N+3}}{\pi^{2N+4}} \sum_{k=2}^{\infty} \frac{1}{(2k-1)^{2N} k^2(k-1)^2}.
\end{aligned}$$

The proof of Theorem 6 is complete. \square

With the evaluations

$$\sum_{k=2}^{\infty} \frac{1}{(2k-1)^4 k(k-1)} = 9 - \frac{\pi^4}{24} - \frac{\pi^2}{2}$$

and

$$\sum_{k=2}^{\infty} \frac{1}{(2k-1)^4 k^2(k-1)^2} = -59 + \frac{13\pi^2}{3} + \frac{\pi^4}{6},$$

the choice $N = 2$ in (4.5) yields

$$\frac{P(x)}{(\pi^2 - 4x^2)^2} < x \sec^2 x - \tan x < \frac{Q(x)}{(\pi^2 - 4x^2)^2}, \quad 0 < x < \frac{1}{2}\pi, \quad (4.7)$$

where

$$\begin{aligned}
P(x) &= \frac{2\pi^4}{3} x^3 + \frac{8\pi^2(\pi^2 - 10)}{15} x^5 + \frac{2(322560 + 1680\pi^4 - 672\pi^6 + 17\pi^8)}{315\pi^4} x^7 \\
&\quad + \frac{16(168 - 17\pi^2)}{315} x^9 + \frac{32(17\pi^8 - 161280)}{315\pi^8} x^{11}
\end{aligned}$$

and

$$\begin{aligned}
Q(x) &= \frac{2\pi^4}{3} x^3 + \frac{32(156 - 6\pi^2 - \pi^4)}{3\pi^2} x^5 + \frac{64(-657 + 37\pi^2 + 3\pi^4)}{3\pi^4} x^7 \\
&\quad + \frac{512(285 - 19\pi^2 - \pi^4)}{3\pi^6} x^9 + \frac{512(-354 + 26\pi^2 + \pi^4)}{3\pi^8} x^{11}.
\end{aligned}$$

The inequality (4.7) is an improvement on the inequality (1.9).

Remark 5. In fact, the lower bound in (4.7) is larger than the one in (1.9), and the upper bound in (4.7) is smaller than the one in (1.9). Hence, the inequality (1.9) holds true. If we write (1.9) as

$$\frac{8\pi^4}{15} - \frac{16\pi^2}{3} < \frac{(x \sec^2 x - \tan x)(\pi^2 - 4x^2)^2 - \frac{2\pi^4}{3} x^3}{x^5} < \frac{256}{\pi^2} - \frac{8\pi^2}{3},$$

we find that

$$\lim_{x \rightarrow 0} \frac{(x \sec^2 x - \tan x)(\pi^2 - 4x^2)^2 - \frac{2\pi^4}{3} x^3}{x^5} = \frac{8\pi^4}{15} - \frac{16\pi^2}{3}$$

and

$$\lim_{x \rightarrow \frac{1}{2}\pi} \frac{(x \sec^2 x - \tan x)(\pi^2 - 4x^2)^2 - \frac{2\pi^4}{3}x^3}{x^5} = \frac{256}{\pi^2} - \frac{8\pi^2}{3}.$$

Hence, the inequality (1.9) holds for $0 < x < \frac{1}{2}\pi$, and the constants $\frac{8\pi^4}{15} - \frac{16\pi^2}{3}$ and $\frac{256}{\pi^2} - \frac{8\pi^2}{3}$ are best possible.

Appendix: A proof of (2.18)

For $N = 0$ in (2.18), we find that

$$\sum_{k=1}^{\infty} \frac{1}{(4k^2 - 1)^2} = \frac{\pi^2 - 8}{16} \quad \text{and} \quad \frac{1}{8}\psi' \left(\frac{1}{2} \right) - \frac{1}{2} = \frac{\pi^2 - 8}{16}.$$

This shows that the formula (2.18) holds for $N = 0$.

Now we assume that the formula (2.18) holds for some $N \in \mathbf{N}_0 := \mathbf{N} \cup \{0\}$. Then, for $N \mapsto N+1$ in (2.18), by using the induction hypothesis and the following relation:

$$\psi'(z+1) = \psi'(z) - \frac{1}{z^2},$$

we have

$$\begin{aligned} \sum_{k=N+2}^{\infty} \frac{1}{(4k^2 - 1)^2} &= \sum_{k=N+1}^{\infty} \frac{1}{(4k^2 - 1)^2} - \frac{1}{(4(N+1)^2 - 1)^2} \\ &= \frac{1}{8}\psi' \left(N + \frac{1}{2} \right) - \frac{N+1}{2(2N+1)^2} - \frac{1}{(4(N+1)^2 - 1)^2} \\ &= \frac{1}{8}\psi' \left(N + \frac{1}{2} \right) - \frac{1}{8(N + \frac{1}{2})^2} - \frac{N+2}{2(2N+3)^2} \\ &= \frac{1}{8}\psi' \left(N + \frac{3}{2} \right) - \frac{N+2}{2(2N+3)^2}. \end{aligned}$$

Thus, by the principle of mathematical induction, the formula (2.18) holds for all $N \in \mathbf{N}_0$.

Acknowledgement

Some computations in this paper were performed using Maple software.

References

- [1] M. Abramowitz, I.A. Stegun (Editors), Handbook of Mathematical Functions with Formulas, Graphs, and Mathematical Tables, Applied Mathematics Series 55, Ninth printing, National Bureau of Standards, Washington, D.C., 1972.
- [2] G. Bach, *Trigonometric inequality*, Amer. Math. Monthly 87 (1980), no. 1, 62.
- [3] A. Baricz, J. Sándor, Extensions of generalized Wilker inequality to Bessel functions, J. Math. Inequal. 2 (2008) 397–406.
- [4] C.-P. Chen, W.-S. Cheung, Inequalities and solution to Oppenheim's problem, Integral Transforms Spec. Funct. 23 (2012), no. 5, 325–336.

- [5] C.-P. Chen, W.-S. Cheung, Sharpness of Wilker and Huygens type inequalities, *J. Inequal. Appl.* 2012 (2012) 72.
- [6] C.-P. Chen, R.B. Paris, Series representations of the remainders in the expansions for certain trigonometric functions and some related inequalities. I (2016).
- [7] C.-P. Chen, F. Qi, A double inequality for remainder of power series of tangent function, *Tamkang J. Math.* 34 (2003), no.4, 351–355.
- [8] C.-P. Chen, J. Sándor, Inequality chains for Wilker, Huygens and Lazarević type inequalities, *J. Math. Inequal.* 8 (2014), no. 1, 55–67.
- [9] H.-F. Ge, New sharp bounds for the Bernoulli numbers and refinement of Becker–Stark inequalities, *J. Appl. Math.* 2012 (2012), Article ID 137507.
- [10] I.S. Gradshteyn, I.M. Ryzhik, *Table of integrals, series, and products*. Translated from the Russian. Sixth edition. Translation edited and with a preface by Alan Jeffrey and Daniel Zwillinger. Academic Press, Inc., San Diego, CA, 2000.
- [11] B.-N. Guo, B.-M. Qiao, F. Qi, W. Li, On new proofs of Wilker inequalities involving trigonometric functions, *Math. Inequal. Appl.* 6 (2003) 19–22.
- [12] C. Huygens, *Oeuvres Completes 1888-1940*, Société Hollondaise des Science, Haga.
- [13] D. S. Mitrinović, *Analytic Inequalities*, Springer-Verlag, Berlin, 1970.
- [14] C. Mortici, The natural approach of Wilker-Cusa-Huygens inequalities, *Math. Inequal. Appl.* 14 (2011) 535–541.
- [15] C. Mortici, A subtly analysis of Wilker inequality, *Appl. Math. Comput.* 231 (2014) 516–520.
- [16] E. Neuman, One- and two-sided inequalities for Jacobian elliptic functions and related results, *Integral Transforms Spec. Funct.* 21 (2010) 399–407.
- [17] E. Neuman, On Wilker and Huygens type inequalities, *Math. Inequal. Appl.* 15 (2012), no. 2, 271–279.
- [18] E. Neuman, J. Sándor, On some inequalities involving trigonometric and hyperbolic functions with emphasis on the Cusa-Huygens, Wilker, and Huygens inequalities, *Math. Inequal. Appl.* 13 (2010) 715–723.
- [19] F.W.J. Olver, D.W. Lozier, R.F. Boisvert, C.W. Clarks (eds.), *NIST Handbook of Mathematical Functions*, Cambridge University Press, New York, 2010.
- [20] M. C. Papenfuss, *Problem E2739*, *Amer. Math. Monthly* **85** (1978), No. 9, 765.
- [21] I. Pinelis, L’Hospital rules of monotonicity and Wilker-Anglesio inequality, *Amer. Math. Monthly*, 111 (2004) 905–909.
- [22] J. S. Sumner, A. A. Jagers, M. Vowe, J. Anglesio, Inequalities involving trigonometric functions, *Amer. Math. Monthly* 98 (1991) 264–267.
- [23] Z. Sun, L. Zhu, Some refinements of inequalities for circular functions, *J. Appl. Math.* 2011, Art. ID 869261.
- [24] J. B. Wilker, *Problem E 3306*, *Amer. Math. Monthly* 96 (1989) 55.
- [25] S.-H. Wu, On extension and refinement of Wilker’s inequality, *Rocky Mountain J. Math.* 39 (2009) 683–687.
- [26] S.-H. Wu, A. Baricz, Generalizations of Mitrinović, Adamović and Lazarević’s inequalities and their applications, *Publ. Math. Debrecen* 75 (2009) 447–458.
- [27] S.-H. Wu, H. M. Srivastava, A weighted and exponential generalization of Wilker’s inequality and its applications, *Integral Transforms and Spec. Funct.* 18 (2007) 529–535.
- [28] S.-H. Wu and H. M. Srivastava, A further refinement of Wilker’s inequality, *Integral Transforms and Spec. Funct.* 19 (2008) 757–765.
- [29] L. Zhang and L. Zhu, A new elementary proof of Wilker’s inequalities, *Math. Inequal. Appl.* 11 (2008) 149–151.
- [30] J.-L. Zhao, Q.-M. Luo, B.-N. Guo, F. Qi, Remarks on inequalities for the tangent function. *Hacet. J. Math. Stat.* 41 (2012), no. 4, 499–506.
- [31] L. Zhu, A new simple proof of Wilker’s inequality, *Math. Inequal. Appl.* 8 (2005) 749–750.
- [32] L. Zhu, Some new inequalities of the Huygens type *Comput. Math. Appl.* 58 (2009) 1180–1182.

- [33] L. Zhu, Some new Wilker-type inequalities for circular and hyperbolic functions, *Abstr. Appl. Anal.* 2009 (2009), Article ID 485842.
- [34] L. Zhu, A source of inequalities for circular functions, *Comput. Math. Appl.* 58 (2009) 1998–2004.
- [35] L. Zhu, Inequalities for Hyperbolic functions and their Applications, *J. Inequal. Appl.* 2010 (2010), Article ID 130821,