

Transparent soil microcosms allow 3D spatial quantification of soil microbiological processes in vivo

Helen F. Downie, Tracy A. Valentine, Wilfred Otten,
Andrew J. Spiers, Lionel X. Dupuy

This is an Accepted Manuscript of an article published in *Plant Signaling & Behavior* online [December 22, 2014], available online: <http://dx.doi.org/10.4161/15592316.2014.970421>

Reproduced by permission of The Society of Plant Signaling and Behavior (<http://www.plantbehavior.org>)

Citation: Downie, H.F., et al. 2014. Transparent soil microcosms allow 3D spatial quantification of soil microbiological processes in vivo. *Plant Signaling & Behavior* 9(10): e970421. Available from DOI: <http://dx.doi.org/10.4161/15592316.2014.970421>

1 ***Short Communication***

2

3 **Transparent soil microcosms allow 3D spatial quantification of soil**
4 **microbiological processes in vivo**

5

6 **Helen F. Downie*^{1,2}, Tracy A. Valentine¹, Wilfred Otten², Andrew J. Spiers², Lionel X.**
7 **Dupuy¹**

8 1. The James Hutton Institute, Invergowrie, Dundee, UK;

9 2. The SIMBIOS Centre, Abertay University, Bell Street, Dundee, UK.

10

11 **Submitted:** 28 May, 2014

12 **Accepted:**

13

14 **Correspondence to:**

15 Helen F. Downie

16 Email: helen.downie@manchester.ac.uk,

17 Current address: Williamson Research Centre for Molecular Environmental Science,

18 University of Manchester, Manchester, UK

19

20 **Abstract**

21 The recently developed transparent soil consists of particles of Nafion, a polymer with a low
22 refractive index (RI), which is prepared by milling and chemical treatment for use as a soil
23 analogue. After the addition of a RI-matched solution, confocal imaging can be carried out in
24 vivo and without destructive sampling. In a previous study, we showed that the new substrate

25 provides a good approximation of plant growth conditions found in natural soils. In this
26 paper, we present further development of the techniques for detailed quantitative analysis of
27 images of root-microbe interactions in situ. Using this system it was possible for the first time
28 to analyse bacterial distribution along the roots and in the bulk substrate in vivo. These
29 findings indicate that the coupling of transparent soil with light microscopy is an important
30 advance towards the discovery of the mechanisms of microbial colonisation of the
31 rhizosphere.

32

33 **TEXT**

34 Plant growth promoting rhizobacteria (PGPR) enhance plant health and yield via complex
35 interactions with the roots and soil ¹⁻³. Rhizobacteria can offer the plant protection from
36 pathogenic microorganisms by outcompeting them and through the promotion of plant
37 growth via the release of plant hormones ⁴. They can also aid plant uptake of nutrients via the
38 rhizosphere, for example by releasing iron-scavenging siderophores ^{4, 5}. The spatial and
39 temporal heterogeneity of soil and the rhizosphere undoubtedly influences the communities
40 and function of bacteria which inhabit niches where nutrients are available in soil ⁶. However,
41 studying the interactions between soil bacteria and their physical habitat is currently very
42 challenging partly due to the lack of conventional laboratory techniques and protocols. Light
43 microscopy cannot be used to observe soil in depth because soil is opaque. X-ray imaging
44 techniques are suitable for studying the soil structure but cannot simultaneously resolve
45 microorganisms ⁷. Although many molecular methods can be used to identify the structure of
46 soil microbial communities ⁸, most do not provide insight into their spatial arrangements. In
47 contrast, recent applications of FISH (fluorescent *in situ* hybridization) have proved
48 successful to analyse spatial distribution of microorganisms in soil, but the method is not
49 suitable to study dynamic processes because samples need to be fixed prior to imaging ⁹.

50 Previously, we published a study describing a new transparent soil analogue for imaging plant
51 roots using optical microscopy [10](#). It consists of a matrix of solid particles of the low
52 refractive index (RI) ionomer, Nafion, water with plant nutrients and air. Transparent soil can
53 be saturated with a RI matched liquid to reveal biological structures within. Further to this
54 work, we have applied transparent soil to the observation of PGPR spatial interactions with
55 roots and soil particles non-destructively, in vivo and in situ. Quantitative analysis methods
56 were developed to study the spatial distribution of PGPR *Pseudomonas fluorescens* SBW25
57 in transparent soil, on the surface of *Lactuca sativa* (lettuce) roots and in the surrounding
58 transparent soil, in relation to the pore geometry. The effect of substrate parameters on the
59 colonisation of roots was also tested by varying the substrate particle size. The aims were to
60 measure the effect of plants and substrate on the abundance of PGPR both on root and on the
61 surrounding particles. After inoculation of the transparent substrate with a culture of GFP-
62 tagged *P. fluorescens*, one day old *L. sativa* seedlings were added to the microcosms. The
63 microcosms were sealed and incubated for 5 days allowing the plants to grow and the
64 bacteria to colonise the roots. The transparency of the substrate allowed images to be
65 captured on a 3D grid using confocal microscopy, thus sampling the microbial abundance at
66 points along the roots and in the bulk soil at 2 distances from the root (supplementary
67 information, figure S1). Fluorescent labelling with a range of fluorophores allowed
68 discrimination of bacteria (GFP), root tissue (calcofluor) and the surfaces of solid Nafion
69 particles (sulphorhodamine-B) (Figure 1), which facilitated image analysis (Figure 2).

70

71 Bacteria were most abundant on the root surfaces, or rhizoplane, and on the surfaces of
72 Nafion particles (Figure 1). Colonisation on the root surface was concentrated in the
73 intercellular junctions of the root epidermal cells (visual observation in 3 samples, e.g. Figure
74 1C), which was similar to observations of field-grown wheat roots [11](#). Watt *et al.* quantified

75 the fraction of the volume of soil occupied by *Pseudomonas* spp. found in wheat
76 rhizospheres. Results showed that on average 15% ¹¹ of the soil volume was occupied by
77 *Pseudomonas* spp. We did not characterise the colonisation of lettuce root by *Pseudomonas*
78 spp. in soil, however, the overall mean rhizosphere volume occupied by *P. fluorescens* in the
79 present study is of the same order of magnitude (10%) as those measured by Watt *et al.*
80 Further studies comparing rhizosphere colonisation with the same plant and bacterial species
81 in both soil and transparent soil would allow a more accurate comparison of the two
82 substrates for this application. Bacterial fluorescence was detected in the pore spaces of the
83 substrate, although at a lower level than on the surfaces (Figures 1, 2A). Image analysis also
84 revealed that the abundance of bacteria in positions with no roots (Figure 2Bi, positions A1-3
85 and B1-3), was constant and independent of image position, particle size and whether a plant
86 was present or not in the chamber. This may indicate that the effect of the plants on soil
87 microbial abundance could be limited to the substrate directly adjacent (i.e. < 1.5 mm) to the
88 root. Along the x axis (horizontal), in samples with plants, the number of discrete bacterial
89 aggregates and the average size of the aggregates was greater on the root (position R) than at
90 1.5 mm (position A) and 3 mm (position B) from the root, and there was no significant
91 difference in bacterial abundance or aggregate number between positions A and B (Figure
92 2Bii). In samples with no plants, there was no difference in bacterial abundance along the X
93 axis (horizontal positions). Along the Y axis (vertical), the number of bacterial aggregates
94 was lower at the root tip (position 1, Figure 2Bii) than the two positions further from the tip
95 (position 2 and 3, Figure 2Bii) but when the percentage area of the image with bacterial
96 fluorescence was used to quantify abundance, there was no difference along the roots (data
97 not shown). In samples with no plants present, the average size of bacterial aggregate was
98 lowest at position 1 and highest at position 3, therefore the points closest to the surface of the
99 substrate had the largest bacterial aggregates (Figure 2Biii). This could be due to a higher

100 concentration of dissolved oxygen closer to the surface, which has been observed in sludge
101 with better bacterial flocculation at high dissolved oxygen concentrations [12](#).
102
103 Several studies have described the distribution of PGPR on the surface of plant roots with a
104 range of, and sometimes contrasting results. High bacterial abundance was found on the root
105 tips [11, 13-15](#) and at root branching zones [13](#). Yet other studies reported an absence or scarcity of
106 bacterial colonisation at the root tips [16-20](#) perhaps caused by the high turnover of mucilage and
107 border cell at the root apex [20](#). It is likely that the choice of the technique used to determine
108 bacterial numbers along the root has a strong influence on bacterial count estimates. Methods
109 based on colony forming units (CFU) are inaccurate because they rely on taking samples and
110 this is difficult on the root tip, and only bacteria that grow well in lab cultures can be quantified.
111 Microscopy techniques such as SEM are usually limited to detect bacteria embedded within
112 the mucilage [16](#), and methods that requires fixing of samples, e.g. FISH, are susceptible to
113 perturbation for example when washing the roots prior to imaging [11](#). The method described
114 in the current study involved the addition and removal of liquids to and from the substrate.
115 Although fluxes of water are common in soil due to rainfall or irrigation, the filling of soil
116 samples by the matching liquid has the potential to induce anaerobic stress in the plant and
117 bacteria over long periods. This effect was minimised by using fresh aerated solutions and by
118 limiting the length of time during which the substrate was saturated. There are numerous
119 non-destructive methods to image in soil, e.g. X-rays, Neutron and Magnetic Resonance
120 Imaging [21-23](#). These do not rely on filling samples in liquid, but the methods are not able to
121 resolve many micro-organisms, and imaging of biological processes such as gene expression
122 or cell division is not possible. Molecular methods are developing rapidly, but currently these
123 are either destructive⁹, or unable to resolve spatial or temporal processes e.g. T-RFLP [24](#).

124

125 The rhizosphere hosts large and diverse bacterial communities that establish sophisticated
126 modes of interactions with plant roots. To date, it has been difficult to characterise such
127 interactions because observation of roots and bacteria in depth and over time has been limiting
128 ²⁵. The model system described here overcomes many previous technological limitations. It
129 combines the ability to grow biological organisms in a physically complex soil-like
130 environment with optical microscopy ²⁵ and to detect multiple fluorescent signals in situ. The
131 application of transparent soil microcosms is not limited to the study of roots and soil bacteria
132 and it holds potential for studying the function of other soil organisms. Future developments
133 could see the introduction of a diversity of microorganisms such as mycorrhizal fungi,
134 nematodes, small invertebrates, or the incorporation of bacterial communities composed of
135 several functional types (e.g. predators and prey). Exploiting this potential now requires
136 exploring, testing and analysing biological activity in transparent soil microcosms to better
137 understand the benefits and limitations of the technology.

138 **Acknowledgements**

139 HFD received funding from a joint studentship from Abertay University and The James Hutton
140 Institute. The James Hutton Institute receives support from the Scottish Government Rural and
141 Environment Science and Analytical Services Division (RESAS, Workpackage 3.3 and 3.4).

142

143 **References**

- 144 1. Urashima Y, Hori K. Selection of PGPR which promotes the growth of spinach. Japanese
145 J Soil Sci Plant Nutr 2003; 74:157-62.
- 146 2. Abbas-Zadeh P, Saleh-Rastin N, Asadi-Rahmani H, Khavazi K, Soltani A, Shoary-Nejati
147 AR, et al. Plant growth-promoting activities of fluorescent pseudomonads, isolated from
148 the Iranian soils. Acta Physiol Plant 2010; 32:281-8.

- 149 3. Dey R, Pal KK, Bhatt DM, Chauhan SM. Growth promotion and yield enhancement of
150 peanut (*Arachis hypogaea* L.) by application of plant growth-promoting rhizobacteria.
151 Microbiol Res 2004; 159:371-94.
- 152 4. Hayat R, Ali S, Amara U, Khalid R, Ahmed I. Soil beneficial bacteria and their role in
153 plant growth promotion: a review. Ann Microbiol 2010; 60:579-98.
- 154 5. Saha R, Saha N, Donofrio RS, Bestervelt LL. Microbial siderophores: a mini review. J
155 Basic Microbiol 2013; 53:303-17.
- 156 6. Vos M, Wolf AB, Jennings SJ, Kowalchuk GA. Micro-scale determinants of bacterial
157 diversity in soil. FEMS Microbiol Rev 2013; 37:936-54.
- 158 7. Fischer D, Pagenkemper S, Nellesen J, Peth S, Horn R, Schloter M. Influence of non-
159 invasive X-ray computed tomography (XRCT) on the microbial community structure and
160 function in soil. J Microbiol Meth 2013; 93:121-3.
- 161 8. Ranjard L, Poly F, Nazaret S. Monitoring complex bacterial communities using culture-
162 independent molecular techniques: application to soil environment. Res Microbiol 2000;
163 151:167-77.
- 164 9. Eickhorst T, Tippkötter R. Improved detection of soil microorganisms using fluorescence
165 in situ hybridization (FISH) and catalyzed reporter deposition (CARD-FISH). Soil Biol
166 Biochem 2008; 40:1883-91.
- 167 10. Downie H, Holden N, Otten W, Spiers AJ, Valentine TA, Dupuy LX. Transparent soil for
168 imaging the rhizosphere. Plos One 2012; 7.
- 169 11. Watt M, Hugenholtz P, White R, Vinnik K. Numbers and locations of native bacteria on
170 field-grown wheat roots quantified by fluorescence in situ hybridization (FISH).
171 Environmental Microbiology 2006; 8:871-84.

- 172 12. Liao BQ, Lin HJ, Langevin SP, Gao WJ, Leppard GG. Effects of temperature and
173 dissolved oxygen on sludge properties and their role in bioflocculation and settling.
174 Water Res 2011; 45:509-20.
- 175 13. Jaeger CH, Lindow SE, Miller W, Clark E, Firestone MK. Mapping of sugar and amino
176 acid availability in soil around roots with bacterial sensors of sucrose and tryptophan.
177 Appl Environ Microbiol 1999; 65:2685-90.
- 178 14. Darwent MJ, Paterson E, McDonald AJS, Tomos AD. Biosensor reporting of root
179 exudation from *Hordeum vulgare* in relation to shoot nitrate concentration. J Exp Bot
180 2003; 54:325-34.
- 181 15. Paterson E, Sim A, Standing D, Dorward M, McDonald AJS. Root exudation from
182 *Hordeum vulgare* in response to localized nitrate supply. J Exp Bot 2006; 57:2413-20.
- 183 16. Lugtenberg BJJ, Dekkers L, Bloemberg GV. Molecular determinants of rhizosphere
184 colonization by *Pseudomonas*. Ann Rev Phytopathol 2001; 39:461-90.
- 185 17. Kragelund L, Nybroe O. Competition between *Pseudomonas fluorescens* Ag1 and
186 *Alcaligenes eutrophus* JMP134 (pJP4) during colonization of barley roots. FEMS
187 Microbiol Ecol 1996; 20:41-51.
- 188 18. Gamalero E, Lingua G, Tombolini R, Avidano L, Pivato B, Berta G. Colonization of
189 tomato root seedling by *Pseudomonas fluorescens* 92rkG5: Spatio-temporal dynamics,
190 localization, organization, viability, and culturability. Microb Ecol 2005; 50:289-97.
- 191 19. Simons M, vanderBij AJ, Brand I, deWeger LA, Wijffelman CA, Lugtenberg BJJ.
192 Gnotobiotic system for studying rhizosphere colonization by plant growth-promoting
193 *Pseudomonas* bacteria. Molecular Plant-Microbe Interactions 1996; 9:600-7.
- 194 20. Humphris SN, Bengough AG, Griffiths BS, Kilham K, Rodger S, Stubbs V, et al. Root
195 cap influences root colonisation by *Pseudomonas fluorescens* SBW25 on maize. FEMS
196 Microbiol Ecol 2005; 54:123-30.

- 197 21. Davey E, Wigand C, Johnson R, Sundberg K, Morris J, Roman CT. Use of computed
198 tomography imaging for quantifying coarse roots, rhizomes, peat, and particle densities
199 in marsh soils. *Ecol Appl* 2011; 21:2156-71.
- 200 22. Pohlmeier A, Oros-Peusquens A, Javaux M, Menzel MI, Vanderborght J, Kaffanke J, et
201 al. Changes in soil water content resulting from *Ricinus* root uptake monitored by
202 Magnetic Resonance Imaging. *Vadose Zone J* 2008; 7:1010-7.
- 203 23. Olson MS, Ford RM, Smith JA, Fernandez EJ. Quantification of bacterial chemotaxis in
204 porous media using Magnetic Resonance Imaging. *Environ Sci Technol* 2004; 38:3864-70.
- 205 24. Gao DW, Tao Y. Current molecular biologic techniques for characterizing environmental
206 microbial community. *Front Env Sci Eng* 2012; 6:82-97.
- 207 25. Yang Z, Downie H, Rozbicki E, Dupuy LX, MacDonald MP. Light Sheet Tomography
208 (LST) for in situ imaging of plant roots. *Opt Express* 2013; 21:16239-47.

209

210 **Figure Legends**

211

212 **Figure 1**

213 Maximum projection confocal images of GFP-labelled *Pseudomonas fluorescens* colonies
214 (green) on the surface of lettuce root tissues (grey) *in situ* in transparent soil with Nafion
215 particles from the substrate labelled with sulphorhodamine B fluorescent dye also visible
216 (red). (A) The majority of the bacterial fluorescens is associated with the root tissue. Scale
217 bar = 150 μ m. (B) Bacteria are present on the root tip and in this case also the surfaces of
218 Nafion particles in close proximity to the root have bacterial fluorescens associated with
219 them. Scale bar = 150 μ m. (C) At higher resolution, bacterial colonisation was predominantly
220 observed in the intercellular junctions of root epithelial cells. Scale bar = 45 μ m.

221

222

223 **Figure 2**

224 Quantification of *Pseudomonas fluorescens* in the rhizosphere. A) Bacteria, Nafion particles
225 and roots were processed sequentially to allow quantification. (i-ii) Bacterial fluorescence
226 before and after processing with a median filter and thresholding facilitated measuring the
227 bacterial abundance. Scale bar = 40 μm . (iii-iv) Original images of particle surfaces were
228 processed and skeletonised. Grey lines in (iv) represent skeleton of particle surfaces in (iii). It
229 was then possible to select the volumes inside particles (shown here in blue) to measure them
230 to correct for available area (pore space). Scale bar = 200 μm . (v-vi) Example image of a
231 section of lettuce root before and after the application of a median filter and subsequent
232 thresholding were applied. This allowed the selection of the internal volume of the root for
233 measurement (shown in blue). Scale bar = 200 μm . B) Quantification of bacterial distribution
234 in transparent soil with small (500 – 850 μm) and large particles (850 – 1200 μm). The
235 positions R1 to B3 represent a 3×3 grid of points on and around the roots, where R is on the
236 root and A and B are at intervals perpendicular to the root. 1 is the root tip and 2 and 3 are
237 closer to the shoot. See Figure S1 for schematic. (i) There was higher bacterial abundance in
238 images that include a section of plant root. At all other positions, there was a consistent area
239 of bacterial fluorescence as a proportion of the area of background in images without plant
240 roots. These values were corrected for available area. (ii) Number and (iii) average size of
241 bacterial aggregates at the 3 horizontal (X) positions (R, A & B) and at the 3 vertical (Y)
242 positions (1, 2 & 3) in samples with or without plants. Letters above the bars indicate the
243 results of Fisher's protected LSD tests.

244

245 **Figure S1**

246 Protocol for the development and imaging of transparent soil microcosms. A) The preparation
247 of the transparent soil material follows three main steps. First, Nafion precursor particles are
248 milled in liquid nitrogen to obtain suitable particle size distribution. In a second stage,
249 particles are treated chemically to give them an anionic charge [11, S1]. Finally, the exchange
250 sites are saturated with the cations from Murashige & Skoog basal plant nutrient medium and
251 the transparent soil medium water content is adjusted for optimal growth conditions. B) The
252 culture and imaging of root and bacteria in transparent soil requires several steps. Seedlings
253 and bacteria are inoculated in the transparent soil substrate at the start of each experiment and
254 grow for 5 days. Samples are then saturated with half-strength M&S medium containing
255 1 mg ml^{-1} fluorescent brightener to stain the root tissue. Immediately before imaging, this
256 solution was removed and replaced with pure Percoll (Sigma-Aldrich Co.) containing $1 \mu\text{g}$
257 ml^{-1} sulphorhodamine B (Sigma-Aldrich Co.). C) Purpose-built transparent containers were
258 constructed for the experiment. Containers were made of a microscope slide and long cover
259 glass with a 4 mm spacer between them on 3 sides and an opening at the top. D) 27 positions
260 were imaged in each sample following a 3 by 3 by 3 regular grid. When plants were present,
261 the origin of the sampling grid is the root tip and the other positions are obtained along and
262 perpendicular to the roots. In the control samples, the reference point R1 was chosen
263 arbitrarily.

264

265